Fax: (947) 886-4655



VITAL CARE OF AUBURN HILLS (0429)

Address: 1532 N Opdyke Road, Suite 700 City/State/Zip: Auburn Hills, Michigan 48326

Essential Data To process the referral, please provide the following minimum information:

- Patient demographic information
- Copy of insurance card (front and back)
- History and physical documentation (including documentation of infection history and diagnosis, recent hospitalizations)
- Recent medication list, IV access information, and home health agency if applicable Currently received and/or prior filed therapies along with lengths of treatment and/or reason for discontinuation
- TB/PPD test if applicable
- Any additional relevant clinical documentation (labs and tests)

Patient Demographics					
Last Name:		First Name:	SSN:		
Date of Birth:	Sex:	Height (cm):	Weight (kg):		
Allergies:					
Address:					
City:		State:	Zip:		
Home Phone:		Mobile Phone:			
Emergency Contact Name:		Emergency Contact Phone:			
□ Crohn's Disease (ICD-10 K50.90)		□ Ulcerative Colitis (ICD-10 K51.90)			
□ Other Diagnosis/ICD-10:		□ Other Diagnosis/ICD-10:			
If unable to find diagnosis/ICD-10 code, please refer to ICD-10-CM (cdc.gov) for reference.					

Patient Insurance Information			
Primary Insurance Name:	Secondary Insurance Name:		
Policy Holder Name:	Policy Holder Name:		
Relation to Policy Holder:	Relation to Policy Holder:		
Member ID:	Member ID:		
Group ID:	Group ID:		

Prescription*			
Medication	Dose/Strength	Directions	
□ Entyvio (vedolizumab)	300mg vial	□ INITIAL: Infuse 300mg IV at week 0, 2, 6, then every 8 weeks thereafter □ MAINTENANCE: Infuse 300mg IV every weeks (max duration of 1 year) □ MAINTENANCE: after at least two IV infusions, administer 108mg SQ every 2 weeks thereafter (max duration of 1 year)	
□ Inflectra (infliximab)□ Remicade (infliximab)□ Renflexis (infliximab)	100mg vial	□ INITIAL: Infusemg/kg IV at week 0, 2, 6, then every 8 weeks thereafter □ MAINTENANCE: Infusemg/kg IV every weeks (max duration of 1 year) □ Other	
□ Stelara (ustekinumab)	130mg vial 90mg (two 45mg vials)	□ INITIAL: Weight based dosing, infuse IV □ 55kg or less: 260mg (2 vials) □ 55kg to 84kg: 390mg (3 vials) □ >85kg: 520mg (4 vials) □ MAINTENANCE: Inject 90mg SC 8 weeks after initial dose, then every 8 weeks thereafter (max duration of 1 year)	
□ Skyrizi (risankizumab)	600mg/10mL vial	□ INITIAL: Infuse 600mg/10mL IV at week 0, 4, and 8 □ MAINTENANCE: Inject 360mg/2.4mL SQ via injector at week 12, then every 8 weeks thereafter (max duration of 1 year)	

Additional Management Orders*			
Hydration Protocol: Per Pharmacy Protocol Infuse 250mL of 0.9% Sodium Chloride Per Provider Protocol	Premedication Protocol Per Pharmacy Protocol Acetaminophen 650mg oral given 30 minutes prior to infusion Diphenhydramine 25mg oral given 30 minutes prior to infusion Per Provider Protocol		
Anaphylaxis Protocol: □ Per Pharmacy Protocol □ Per Provider Protocol	Flush Protocol: □ Per Pharmacy Protocol Flush 10mL of 0.9% Sodium Chloride pre and post infusion □ Per Provider Protocol		
Labs Protocol: Per Pharmacy Protocol Per Provider Protocol	Other:		
*Product selection permitted unless dispense as written is checked or clearly written on order form. *Vials will be rounded to the nearest vial if appropriate unless clearly written to dispense dose as written from provider. *If order is not specified, it will be considered as a one-time order and must be rewritten.			

Notes to Pharmacy

	First [Dose to	be c	given	in	home	
_	Chille	d Nivea	:	.:-:			_

□ Skilled Nursing visits as required

Referring Provider Information			
Ordering Provider's Name:			
NPI:	Specialty:		
Facility Name:			
Address:			
City:	State:	Zip:	
Office Phone:	Office Fax:		
Dispense as Written			
Provider Signature:		Date:	

Lauthorize Vital Care Infusion Services LLC and its representatives to initiate any insurance prior authorization process that is required for this prescription and for any future refills of the same prescription for the patient listed above which I order. I understand that I can revoke this designation at any time by